

copy

well  
Dnc.

onal  
king  
pro-  
ties  
dif-  
hus,  
ent,  
CD  
ther  
zed  
ght-  
mo-  
nts,  
um.  
for

the  
wer  
nce

tion  
- S<sub>0</sub>  
S n<sub>0</sub>  
f(v)  
)=

Microscopy

initially

(8-33)

the con-  
-time of

(8-34)

l absorp-  
Equation  
the more  
n 8-34 is  
n is sub-  
Equation  
ansitions.  
y as long  
ited state  
h of light  
tially can

script

mon

slide

polar

sec<sup>-1</sup>.

nally

triplet

er by

The

ence.

from

mall.

ctro-

ribed

triplet

onds

ermal

us. in

rigid

order

ather

ence

inglet

n de-

ed by

(8-38)

(8-39)

ence

(8-40)

(8-41)

py  
to is he ds he ant ce pre me he

of er ds ed an nd nd nt u- at ay- D- re he of ed ver ase us

a- lly

intensity

length  $\lambda$  is  
ability of  
length  $\lambda$ ;  
vector.  
absorption

(8-42)

$l$  and  $d$   
position of  
excited  
how the

relative  
(0.70), or  
cent. The  
d for the

on 8-41.  
excite the  
ission as  
pulse will  
singlets

(8-43)

equations  
directly,  
shows a  
e single

nt. Now  
choice of

... ..

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]


[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]



H<sub>2</sub>

ds of  
vent-  
rans-  
reous  
rigid  
ly be  
on a  
ound  
cently  
um is

530

lives  
d by

seek  
aid.

nee  
ons  
pre-

more  
plet.

ter-

Any  
the

n is  
ular

how  
n of

of a  
plex

more  
ough

f the  
pre-

and





-singlet energy-  
Donor (colored line)  
(black line) and  
of  $v^{-1}$  is shaded.

it ( $A_n$ ) and donor singlet  
n rapidly converts these  
an efficient process, the  
transfer resonance shifts the  
donor becomes quenched.  
resc. This process, from  
note that the two chromo-  
Energy transfer will take  
the concentration is high  
A or so.

and acceptor excitation  
non-interaction that leads to  
probational relaxation (vary-  
dly and tends to prevent a  
action that would relax the  
ate the donor.

Microscopy

(8-53)

portant

is that  
fore, in  
which  
English-  
donor  
ected  
It can  
0-3 M.

iciency  
done  
rate

(8-54)

sixth  
R-3

(8-55)

(8-56)

osorp-  
efined  
eptor:  
mplex  
both  
scence  
ties in

lum yield of the donor.

occurs over a band of the donor occurs over a frequency  $\nu$ . The total yield at each frequency:

$\eta^2$

or numerical constants from the fact that  $\bar{\nu}$  as interaction potential at quantity  $\kappa^2$ . Removing we did above implies  $\eta^2$ . This may not always be true. We know that polarization can be a  $\kappa^2$  value still are a factor. Fortunately, the equation 8-57 in this form:

to use spectra measured equations 8-55 and 8-56

a function of  
-naphthyl  
 $n = 1$  to  $12$ . The  
Equation 8-57.  
England, *Proc.*  
[1967].]

the  $y$  axis. The  
lled light are  
 $x$  axis. There  
long this axis

ed to a frozen  
ted molecules  
nce time scale  
on transition  
y coordinates.  
al to  $(\mu \cdot E)^2$ .  
y is the angle  
dipoles near  
selection.  
probability of  
ne probability  
convenient to  
ve number of  
is  $\sin \theta \, d\theta \, d\phi$ .  
olecules per-  
g a molecule  
Therefore, the

ced by  
propagating  
is pro-  
ceived  
particular

$\psi(\theta, \phi)$ , and

3,5 (8-61)

1 to  $|\mu \cdot \hat{r}|^2$ ,  
 $\sin^2 \theta \cos^2 \phi$ ;  
probabilities


$\sin^2 \theta = 1, 5$   
(8-62)

convenient  
are defined



(8-63)

as you can  
8-63. These  
under any



The results for

(8-70)

(8-71)

ential, each of  
equations 8-70  
copy. At the

$1/2$ ] (8-72)

(8-73)

4-64,  
molecule is a  
the-hydrated  
Equation 8-73  
fluorescence aniso-  
more slowly.  
pilotyl chymo-  
typical value

Equation 10-10  
protein is about  
be  $M/N_0 =$   
centipoise:  $T$   
Equation 8-73 as

12 sec (8-74)

if the protein  
for globular  
weight is 25,000  
predicted and  
elongated shapes

(8-77)

macro-

case,

(8-78)

(8-79)

same

ations

(8-80)

(8-81)

time  
is we  
. The  
as a  
rding  
olec-  
The  
8-24  
ropy  
ional  
mea-  
tein.  
lead,  
much



py versus  
ample in which  
the attach-  
by a macro-  
near portion  
apparent  
r than the  
n these values  
es over which  
relative to the

cular motion  
orientations.  
y will reflect  
k to  $T$  if  $= 0$   
s covered by  
tionary. The  
an enhanced  
linear Perrin  
s can provide  
r of a whole  
quite difficult  
ation.  
can be used  
acid contains  
cited. energy  
pres prior to  
nal absorber.  
polarization  
the chromo-  
rption, these  
r, leading to  
substantially  
e an effective  
ansfer effects  
ze and shape